

Simulations of extreme dry and wet rainy seasons of West Africa and their trend in future climates using a simple vegetation model within ECHAM5

Tim Brücher and Andreas H. Fink

Institute f. Geophysik and Meteorology, University of Cologne, Germany
Centre for Applied Computer Sciences, University of Cologne, Germany



outlook

- Motivation / IMPETUS
- IMPETUS Model Chain (from global to local scale)
- SVege - Simple Vegetation Model
- results
 - present day climate
 - future trends
- conclusion






outlook

Motivation

Rainfall Variability in Tropical West Africa (1921-2007)

$$Index = \frac{\sum_{JJAS} RR_i - CLIM_{JJAS}}{\sigma_{JJAS}}$$

Base period: 1950-1990



Figures: courtesy of A. Fink and S. Kotthaus, IGM, University of Cologne, Germany




Motivation

Rainfall Variability in Tropical West Africa (1921-2007)

$$Index = \frac{\sum_{JJAS} RR_i - CLIM_{JJAS}}{\sigma_{JJAS}}$$

Base period: 1950-1990





Motivation


Rainfall Variability in Tropical West Africa (1921-2007)

$$Index = \frac{\sum_{JJAS} RR_i - CLIM_{JJAS}}{\sigma_{JJAS}}$$

Base period: 1950-1990



what about ...
... future trends
... socioeconomic impacts
... food security
...



- **An Integrated Approach to the Efficient Management of Scarce Water Resources in West Africa**
- **The Catchments of the Ouémé-River (Benin) and the Wadi Drâa (Morocco) are investigated**
- **Relations to the climates of Europe might exist via complex atmosphere-ocean interactions in the area of the tropical/ subtropical and north Atlantic ocean**
- **West Africa: most pronounced inter-decadal variability of climate**
- **for more details see: <http://www.impetus.uni-koeln.de>**



Challenges

concerning future climate trends

- after the IPCC 4AR there is no clear projection for West African precipitation trends (increase / decrease)
 - ➡ it is necessary to put some light on it
- what about changes in the intensity and variability of the West African Monsoon on different time scales
- what about Monsoon onset and brakes in future climate

Project Plan

- **1st Phase:** Data Acquisition and Modeling
(2000-2003)
- **2nd Phase:** Development of Scenarios and Problem Clusters (2003-2006)
- **3rd Phase:** Transfer and Application: Capacity Building and Spatial Decision Support Systems (2006-2009)


Project Plan

- **1st Phase:** Data Acquisition and Modeling
(2000-2003)
- **2nd Phase:** Development of Scenarios and Problem Clusters (2003-2006)
- **3rd Phase:** Transfer and Application: Capacity Building and Spatial Decision Support Systems
(2006-2009)

IMPETUS model chain

- model chain for a statistic - dynamical down-scaling of global projections down to key-region
- starting from 250 km down to 3 km


consider as many aspects as possible to get best boundary conditions for nested models



IMPETUS model chain


- model chain for a statistic - dynamical down-scaling of global projections down to key-region
- starting from 250 km down to 3 km

consider as many aspects as possible to get best boundary conditions for ne




➡ include biosphere: **Simple Vegetation model (SVege)**

Motivation: using SVege within E4



Base period for anomaly: 1951-1994


Motivation: using SVege within E4




Schnitzler et al. (2001)

Base period for anomaly: 1951-1994

SVege: Simple Vegetation Model




SVege: Simple Vegetation Model



SVege includes **no** dependencies of vegetation growth upon temperature and duration of sunshine:

- Implemented for **non-boreal** areas (40°N to 40°S)
- Outside this domain the albedo values from the **Land Surface Parameters** - dataset (LSP, Steffan Hagemann 2002) are used (climatological mean)
- No other changes in land surface parameters are simulated (e. g. roughness length)

SVege: Simple Vegetation Model



SVege includes **no** dependencies of vegetation growth upon temperature and duration of sunshine:

- Implemented for **non-boreal** areas (40°N to 40°S)


our update:

only grid boxes with a vegetation cover higher than 15% are considered

simulated (e. g. roughness length)


SVege: „Cologne version“

standard albedo of ECHAM5




SVege: „Cologne version“

standard albedo of ECHAM5




standard version of SVege




SVege: „Cologne version“


standard albedo of ECHAM5



Cologne Version




Annual cycle of albedo




reproduction of albedo's annual cycle

observed mean June to August precipitation (1960-1999)



Differences in summer precipitation amounts (JJA)



observations – E5 SVege



Differences in summer precipitation amounts (JJA)

observations – E5 SVege



observations – E5 original



Differences in summer precipitation amounts (JJA)

observations – E5 SVege

observations – E5 original




- both versions comprise same strengths and deficiencies
- only small, insignificant differences between the model



-500 -300 -200 -100 0 100 200 300 500 [mm]

-500 -300 -200 -100 0 100 200 300 500 [mm]

decadal variability




decadal variability




significant changes to an higher correlation (up to 0.2) between model and observation using the vegetation model


precipitation annual cycle



precipitation annual cycle




precipitation annual cycle




about extreme dry and wet years

Central Sahel




about extreme dry and wet years


Central Sahel



Guinea Coast



differences between both versions in extreme dry and wet years




Central Sahel E5+SVege: ■ wet years ■ dry years
 E5 original: ■ wet years ■ dry years

differences between both versions in extreme dry and wet years

Tmin

Tmax




Central Sahel E5+SVege: ■ wet years ■ dry years
 E5 original: ■ wet years ■ dry years

differences between both versions in extreme dry and wet years

bowen ratio β

cloud cover




Central Sahel E5+SVege: ■ wet years ■ dry years
 E5 original: ■ wet years ■ dry years



what about future trends ?


increasing variability (AIB)



- increasing standard deviation of the precipitation index
- increasing number of extreme seasons in the Sahel + Guinea Coast (wet and dry)
- decreases are projected for the Sahel using the BI-scenario

significance of changes (AIB)

phase diagram to show variability of standard deviation and mean values




conclusion

- implementation of biosphere to ECHAM5 using a simple vegetation approach (SVege)
 - ➡ small, sometimes insignificant changes to standard version
 - ➡ improved representation of decadal variability
 - ➡ getting same results by a more physical based way
- the simple vegetation approach could be *to simple* for the sophisticated version of ECHAM5 ?
- COSMOS-Earth System Model (ESM); MPI-Hamburg, available in August / September (?)


ideas for future work

- „testing“ the model in comparison to other WAMME experiments
- improvement by using another aerosol climatology (A. Tompkins)

Tanre (1984)



Tegen (1997)



5d forecast for windspeed at 700 hPa



thanks for your interest